

Euclidean 4-simplices and invariants of four-dimensional manifolds: III. Moves $1 \leftrightarrow 5$ and related structures

I. G. Korepanov

Southern Ural State University
76 Lenin avenue
454080 Chelyabinsk, Russia
E-mail: kig@susu.ac.ru

Abstract

We conclude the construction of the algebraic complex, consisting of spaces of differentials of Euclidean metric values, for four-dimensional piecewise-linear manifolds. Assuming that the complex is acyclic, we investigate how its torsion changes under rebuildings of the manifold triangulation. First, we write out formulas for moves $3 \rightarrow 3$ and $2 \leftrightarrow 4$ based on the results of our two previous works, and then we study in detail moves $1 \leftrightarrow 5$. On this basis, we obtain the formula for a four-dimensional manifold invariant.

As an example, we present a detailed calculation of our invariant for sphere S^4 ; in particular, the complex turns out, indeed, to be acyclic.

1 Introduction

This work is third in the series of papers started with papers [1] and [2]. We also use Roman numerals I and II, respectively, for references to those works; in particular, equation (II.1) is the equation (1) from paper [2], or “from paper II”.

Our goal is to construct and investigate a new type of acyclic complexes, wherefrom we should be able to extract invariants of four-dimensional piecewise-linear manifolds. Note, however, that at the time when paper I was being written, it was not yet clear that acyclic complexes were exactly those structures that stood behind the considered algebraic formulas, as well as behind the invariants of *three-dimensional* manifolds from papers [3] and [4]. These complexes were written out in paper II — the whole complex for the three-dimensional case and a part of it — for the four-dimensional case. Concerning the four-dimensional case, it turned out that, in order to construct a value invariant under Pachner moves (i.e., rebuildings of the manifold triangulation) of type $3 \rightarrow 3$ *only*, it is enough to consider only a small central part of the complex

(two vector spaces and one linear mapping between them), whereas the addition of rebuildings $2 \leftrightarrow 4$ brings into consideration one more space (namely, the space of edge deviations) and one more mapping.

Paper I was devoted to moves $3 \rightarrow 3$, and paper II — to moves $2 \leftrightarrow 4$. Therefore, the complex *in full* was not necessary in those papers, and its full form was only announced in paper II (formulas (II.6) and (II.7)).

The contents of the present “paper III” by sections is as follows. In Section 2 we revisit, once again, the dimensionality 3 and write out the acyclic complex from Section 2 of paper II in a slightly different (longer) form, which clarifies the structure of one of the involved spaces as a *factor* space. This simple improvement may simplify considerably our constructions when we pass to higher dimensions.

In Section 3 we present the space of vertex deviations, as we promised in paper II, and its mapping into the space of edge deviations. This enables us to write out in Section 4 our complex for the four-dimensional case in full; we also “make it longer” with respect to paper II in analogy with the three-dimensional case. We still have to prove, though, that our sequence of spaces and mappings is really a complex, and we do that in Section 5.

We would like our constructed complex to be acyclic (i.e., an exact sequence) which would enable us to get manifold invariants out of its torsion. We do not give the full proof of the acyclicity in the general case, limiting ourselves to some remarks in the end of Section 5 (see also a concrete example in Section 9). Then we show that, *assuming the exactness*, the torsion of our complex does not change under Pachner moves (and the exactness itself is conserved under those moves). To this end, we study in Section 6 how the complex changes under moves $3 \rightarrow 3$ and $2 \leftrightarrow 4$, and in Section 7 — under moves $1 \leftrightarrow 5$ (one can say that it was the need to study the moves $1 \leftrightarrow 5$ that required the detailed consideration of the *whole* algebraic complex). In Section 8 we present the final formula giving the invariant of a four-dimensional piecewise-linear manifold (i.e., the invariant of *all* Pachner moves) in terms of the torsion of the complex and other Euclidean geometric values. In Section 9 we show how our machinery works for the sphere S^4 (and check explicitly that, at least for the sphere, our complex is indeed acyclic).

In the concluding Section 10 we discuss the obtained results.

2 Revisiting three dimensions once again: a longer form of the acyclic complex

We recall that the acyclic complex from Section 2 of paper II, corresponding to a triangulation of a three-dimensional manifold M , had the following form:

$$0 \leftarrow (\cdots) \xleftarrow{B^T} (d\omega) \xleftarrow{A} (dl) \xleftarrow{B} (dx \text{ and } dg) \leftarrow 0. \quad (\text{II.1})$$

Here $(d\omega)$ is the space of column vectors made of infinitesimal deficit angles at all edges of the complex, (dl) is the space of columns of differentials of edge lengths, (dx) is

the space of columns of differentials of Euclidean coordinates of the edges of the complex (to be exact, of their lift-ups onto the universal cover), (dg) is the space of columns of differentials of continuous parameters on which the representation $f: \pi_1(M) \rightarrow E_3$ may depend. In particular, (dg) vanishes for manifolds with a finite fundamental group.

We thus continue to use the notations in the style of paper II for linear spaces entering in algebraic complexes, hoping that the convenience of such notations pays for their certain looseness. Of course, from the formal standpoint, for instance, space (dl) is the *tangent space* to the manifold consisting of all sets of positive numbers — “lengths” — put in correspondence to the edges of our triangulation; $(dx$ and $dg)$ (the space of columns of all dx and all dg) is the *direct sum* $(dx) \oplus (dg)$.

Recall that (dx) in sequence (II.1) is the space of differentials of coordinates taken up to those infinitesimal motions of the Euclidean space $\mathbb{R}^3$ which are compatible with the given representation f , i.e. motions commuting with its image $\text{Im } f = f(\pi_1(M))$. Such motions form a subalgebra in the Lie algebra $\mathfrak{e}_3$, which we denote $\mathfrak{a}$. Column (dx) is described explicitly for various cases in Section 2 of paper II. In the three-dimensional case, such description can be made easily, but difficulties may increase when we pass to higher dimensions. That is the reason for rewriting sequence (II.1) in the following, more elegant form.

We now permit ourselves to *change notations* and understand below by (dx) the space of columns of differentials of Euclidean coordinates of the vertices in the complex, with no further conditions, that is, columns of values $(dx_1, dy_1, dz_1, \dots, dx_N, dy_N, dz_N)$. Then our *former* space (dx) is written as the factor $(dx)/\mathfrak{a}$. This suggests an idea of adding one more term to the right of $(dx$ and $dg)$ in sequence (II.1) (and a symmetric term in the left-hand part of the sequence — see below), understanding now (dx) in the new sense:

$$0 \leftarrow (\dots) \leftarrow (\dots) \xleftarrow{B^T} (d\omega) \xleftarrow{A} (dl) \xleftarrow{B} (dx \text{ and } dg) \leftarrow \mathfrak{a} \leftarrow 0. \quad (1)$$

Recall that the whole sequence (this applies to both (II.1) and (1)) is symmetric in the following sense. Each term in it is considered as a vector space with a fixed basis; mappings between them are identified with matrices; and any two matrices at equal distances from the left-hand and right-hand ends of the sequence must be obtained from one another by matrix transposing. In particular, matrix A that gives the mapping $(dl) \rightarrow (d\omega)$ is symmetric.

As for the notation “ $(\dots)$ ”, we are using it for *different* linear spaces whose specific geometrical sense we are not going to investigate (at this moment; still, we know the matrices of mappings between such spaces, for instance, from the symmetry described in the previous paragraph).

The torsion of an acyclic complex is the product of certain minors in the matrices of its linear mappings, taken in alternating degrees ± 1 . If we adopt a convention that the $+$ sign corresponds to the first nontrivial mapping (coming after the zero injection, i.e., for example, in sequence (II.1) that is the second arrow from the right), then the torsion will change to its inverse when we pass from (II.1) to (1). To avoid this, we can *agree* to take the $+$ sign for mapping B in sequence (1), as before. Then *the torsion*

of complex (1) coincides with the torsion of complex (II.1) if we choose a basis in Lie algebra $\mathfrak{e}_3$ in a natural way (namely, three infinitesimal translations along mutually orthogonal axes and three rotations around these very axes).

The full proof of this statement must involve the analysis of all the particular cases from Section 2 of paper II. Here, we will limit ourselves to the case of a lens space $L(p, q)$, with a nontrivial homomorphism $f: \pi_1(L(p, q)) \rightarrow E_3$ whose image consists of rotations around the z axis. To compose the minor of the matrix of mapping $\mathfrak{a} \rightarrow (dx)$, we choose two basis elements in space (dx) , namely, dx_1 and dz_1 , where subscript 1 corresponds to some vertex in the triangulation which we agree to call the “first” one. Now a simple straightforward calculation shows that, indeed, the torsions of complexes (1) and (II.1) coincide, if we choose the basis of the “old” (dx) in the latter complex according to formula (II.2) (it makes sense to remind here that we consider the torsion to within its sign).

3 Vertex deviations and their mapping into edge deviations

We have already used Euclidean coordinates in our constructions a few times, namely, the coordinates of a vertex in the complex were necessary to define the mappings $(dx) \rightarrow (dl)$ in the three-dimensional and four-dimensional cases, and the coordinates of the vector of edge deviation — for the mapping $(d\vec{v}) \rightarrow (dS)$. We can remark that we did not need any connection between the coordinate systems for *different* vertices and/or edges while doing these constructions. Any individual coordinate system could be chosen arbitrarily, for instance, by fixing some angles between coordinate axes and adjacent edges.

We cannot work with no such coordinate systems at all if we want to *fix the bases* in all vector spaces entering in a complex; on the other hand, the torsion of our complexes, as one can check, does not depend on the choice of those systems.

In this Section we, first, define the *vertex deviation* as a tensor value which needs, for its components to be fixed, a coordinate system corresponding to the *vertex*. Next, it will be convenient for us to define how it generates edge deviations for the edges abutting on the given vertex, using the *same* (i.e. corresponding to the vertex) coordinate system for those edges. We *imply* that the components of each edge’s deviation are then transformed into *its own* coordinate system by using a proper orthogonal transformation.

So, we call vertex deviation a bivector (antisymmetric tensor) $d\sigma_{\alpha\beta}$, where $\alpha, \beta = 1, \dots, 4$. Let AB be one of the edges abutting on vertex A . This edge can be also considered as a four-dimensional vector $\overrightarrow{AB}$, whose coordinates we denote as l_α . By definition, a deviation of vertex A equal to $d\sigma_{\alpha\beta}$ generates the deviation of edge $d\vec{v}_{AB}$ with components

$$(d\vec{v}_{AB})_\beta = \frac{1}{L_{AB}} \sum_{\alpha} l_{\alpha} d\sigma_{\alpha\beta}, \quad (2)$$

where $L_{AB} = \sum_{\alpha} l_{\alpha}^2$ is the squared length of edge AB .

In general, $d\vec{v}_{AB}$ is the sum of expression (2) and a similar expression involving the deviation of vertex B (where, of course, vector $\vec{BA}$ must be used instead of $\vec{AB}$). The reasonableness of definition (2) will be clear in Section 5, where we will prove, in particular, that the edge deviations of type (2) generate zero differentials (dS) of two-dimensional face areas. This will be part of the statement that “composition of two neighboring arrows (i.e. linear mappings) in the sequence is zero”, which will justify the name “complex” for that sequence. Let us now pass on to the construction of our sequence in its full form.

4 The full sequence of spaces and mappings in the four-dimensional case

We will write out two “conjugate” sequences, as we did in Section 3 of paper II. Compared with formulas (II.6) and (II.7), we will increase them in length both on the right and on the left, in analogy with the three-dimensional case (Section 2 of the present work), because both the leftmost nonzero space in formula (II.7) and the rightmost one in (II.6) can be represented naturally as factor spaces.

As for the first of mentioned spaces, denoted as $(dx$ and $dg)$ in formula (II.7), here our elongation goes, in principle, the same way as in three dimensions. Thus we pass at once to the second one, denoted $(d\sigma)$ in formula (II.6), i.e. to the space of vertex deviations introduced in the previous Section. It turns out that there exists an easily defined space of “trivial” deviations generating zero edge deviations $d\vec{v}_a$, and it finds its natural place in our complex.

Consider formula (2). We recall that, in it, the tensor value $d\sigma_{\alpha\beta}$ pertains to point A , and l_{α} are the components of vector $\vec{AB}$. If we take into account $d\sigma_{\alpha\beta}$ in point B as well, we get

$$(d\vec{v}_{AB})_{\beta} = \frac{1}{L_{AB}} \sum_{\alpha} l_{\alpha} ((d\sigma_A)_{\alpha\beta} - (d\sigma_B)_{\alpha\beta}). \quad (3)$$

One can see from here that $d\vec{v}_{AB} = 0$ if the difference $(d\sigma_A)_{\alpha\beta} - (d\sigma_B)_{\alpha\beta}$ has the form $\sum_{\gamma,\delta} \epsilon_{\alpha\beta\gamma\delta} l_{\gamma} d\vec{s}_{\delta}$, where $\epsilon_{\alpha\beta\gamma\delta}$ is the totally antisymmetric tensor, $\epsilon_{1234} = 1$, and $d\vec{s}$ is any infinitesimal vector.

Passing on from considering one edge AB to considering all edges in the complex, we take, first, the case of trivial representation $f: \pi_1(M) \rightarrow E_4$ of the fundamental group of our manifold M into the group of motions of the four-dimensional Euclidean space, i.e. the case $\text{Im } f = \{e\}$. This means that all inverse images of any given vertex in the complex get into one and the same point of space $\mathbb{R}^4$ (see Section 3 of paper I), that is, simply speaking, to each vertex A its radius vector $\vec{r}_A$ corresponds unambiguously with components $(r_A)_{\alpha}$ in some Cartesian coordinate system common for the whole complex.

Choose some infinitesimal antisymmetric tensor $d\tau_{\alpha\beta}$ and vector ds_α , and set for each vertex A

$$(d\sigma_A)_{\alpha\beta} = d\tau_{\alpha\beta} + \epsilon_{\alpha\beta\gamma\delta}(r_A)_\gamma ds_\delta. \quad (4)$$

It follows from the foregoing that such set of vertex deviations yields zero deviations for all edges. Note that such columns $d\sigma$ make up a ten-dimensional linear space.

In case representation f is not trivial, any vertex has more than one inverse images in the universal covering, and these inverse images are placed in different points of space $\mathbb{R}^4$. The requirement that formula (4) must give equal results for all such inverse images (after transforming them into the coordinate system corresponding to the given vertex — see Section 3), leads to linear restrictions on admissible $d\tau_{\alpha\beta}$ and ds_β . In the present paper, we do not write out explicitly these restrictions: it will suffice for us to begin with studying just simply connected manifolds, for which $\pi_1(M) = \{e\}$. Nevertheless, we introduce notation $(d\sigma)_0$ for the subspace of columns of those vertex deviations which are correctly determined by formula (4), for *any* manifold M .

Now we are ready to rewrite sequences (II.6) as (II.7) in the renovated (longer) form. Despite the fact that we are still using notation “ $(\dots)$ ” for vector spaces with whose geometric sense we are not concerned now, we have *given the definitions* to all *matrices* of mappings denoted by arrows. So, here are our mutually conjugate sequences:

$$0 \leftarrow (\dots) \leftarrow (\dots) \leftarrow (d\Omega_a) \xleftarrow{(\partial\Omega_a/\partial S_i)} (dS_i) \leftarrow (d\vec{v}_a) \leftarrow (d\sigma) \leftarrow (d\sigma)_0 \leftarrow 0, \quad (5)$$

$$0 \rightarrow \mathfrak{a} \rightarrow (dx \text{ and } dg) \rightarrow (dL_a) \xrightarrow{(\partial\omega_i/\partial L_a)} (d\omega_i) \rightarrow (\dots) \rightarrow (\dots) \rightarrow (\dots) \rightarrow 0. \quad (6)$$

Here, of course, $\mathfrak{a}$ is a subalgebra of Lie algebra $\mathfrak{e}_4$ of motions of Euclidean space $\mathbb{R}^4$. Namely, $\mathfrak{a}$ consists of those motions commuting with the image of group $\pi_1(M)$ in E_4 , in full analogy with the three-dimensional case from Section 2.

5 The sequence is a complex

Now we will show that the composition of any two successive arrows in sequence (5) or, equivalently, (6) equals zero. Thus, we will justify the name “complex” for each of these sequences.

We start with the two arrows of sequence (5) adjacent to the term $(d\sigma)$. It follows directly from formulas (3) and (4) of the previous Section that their composition is zero.

Moving to the left, we consider two arrows around the term $(d\vec{v}_a)$. Consider a triangle ABC — one of the two-dimensional faces of our simplicial complex — and check that the deviations of its edges, if they are given by formulas of type (3), lead to the zero area differential dS_{ABC} .

It is enough to consider the case where only vertex A has a nonzero deviation $d\sigma$. According to formula (2), the lengths of vectors $d\vec{v}$ in Figure 1 are inversely proportional to the triangle’s sides to which they belong. Thus, one can easily deduce that the area of




Figure 1:

triangle $A'B'C'$ multiplied by four *coincides* with the area of triangle ABC and hence $dS_{ABC} = 0$ according to formula (II.8).

Moving further to the left along sequence (5), we must consider two arrows adjacent to the term (dS_i) — but the vanishing of their product has been already proven in Section 5 of paper II.

To investigate the two remaining pairs of arrows, we switch to sequence (6). In terms of that sequence, these are the pairs of arrows around the terms $(dx$ and $dg)$ and (dL_a) . Now it remains to remark that the statements we need can be proved in full analogy to how it was done for the three-dimensional case in Section 2 of paper II and Section 2 of the present work. Moreover, there is even *exactness* in those terms.

As for the exactness in other terms, nothing is known about it as yet in the general case, save that it is clear from the construction that the exactness holds in the left- and rightmost terms $\mathfrak{a}$ and $(d\sigma)_0$ which appeared when we “made longer” our complex. Still, we will see in the following Sections 6 and 7 that if the sequence is exact, then this property is preserved under Pachner moves, that is, exactness does not depend on a triangulation. Besides, the example of sphere S^4 studied below in Section 9 shows that at least for the sphere the sequence *is* exact (i.e., is an acyclic complex).

6 How the algebraic complex changes under moves $3 \rightarrow 3$ and $2 \leftrightarrow 4$

We already know from papers I and II what happens under moves $3 \rightarrow 3$ and $2 \leftrightarrow 4$. Our task in this Section is to reformulate these results while holding strictly to the algebraic language of acyclic complexes.

6.1 Moves $3 \rightarrow 3$

In paper I, devoted to moves $3 \rightarrow 3$, we were considering matrix $(\partial\Omega_a/\partial S_i)$ and its conjugate $(\partial\omega_i/\partial L_a)$. From our current viewpoint, they form the central part of sequence (5) and its conjugate (6). Some of the results of paper I (namely, Theorem 4) do not deal with moves $3 \rightarrow 3$ as such but only reproduce (by somewhat amateurish means) a part of statement that the torsion of a complex does not depend on a specific choice of minors through which it is expressed. It is Theorem 3 of paper I that deals with moves $3 \rightarrow 3$ proper. It seems worthwhile to recall it here once again (with only a very slight reformulation):

Choose in matrix $(\partial\omega_i/\partial L_a)$ a largest square submatrix $\mathcal{B}$ with nonzero determinant. Let $\mathcal{B}$ contain a row corresponding to such face $i = ABC$ that belongs to exactly three 4-simplices. Then those latter can be replaced by three new 4-simplices by a move $3 \rightarrow 3$. After such a replacement, take in the new matrix $(\partial\omega_i/\partial L_a)$ the new submatrix $\mathcal{B}$ containing the same rows and columns, with the only following change: the row corresponding to the face ABC is replaced by the row corresponding to the new face DEF .

The expression

$$\frac{\det \mathcal{B} \cdot \prod_{\substack{\text{over all} \\ \text{4-simplices}}} V}{\prod_{\substack{\text{over all} \\ \text{2-dim. faces}}} S} \quad (7)$$

does not change under such a rebuilding.

We add here the following. The torsion of complex (6) is the alternated product of minors one of which is exactly $\det \mathcal{B}$. With the presented realization of move $3 \rightarrow 3$, *all other minors obviously remain unchanged*. Thus we can say that formula (7) describes the behaviour not only of $\det \mathcal{B}$ but of the whole torsion τ under a move $3 \rightarrow 3$. It is evident also that the acyclicity property as such, if the complex possessed it, is preserved.

6.2 Moves $2 \leftrightarrow 4$

For convenience, we speak about moves $2 \rightarrow 4$, having in mind the obvious invertibility of the following reasoning and formulas to the case of moves $4 \rightarrow 2$.

Under a move $2 \rightarrow 4$, one edge and four two-dimensional faces are added to the simplicial complex. As for the algebraic complex (5), the dimensionality of space of column vectors (dS_i) increases by 4, the dimensionality of space $(d\vec{v}_a)$ — by 3 and the dimensionality of space $(d\Omega_a)$ — by 1. Hence, we can increase by 3 the sizes of the minor of matrix $(\partial S_i/\partial \vec{v}_a)$ and by 1 — those of the minor of matrix $(\partial\Omega_a/\partial S_i)$, and *keep unchanged the minors* of which the torsion is made up *beyond the fragment* $(d\Omega_a) \leftarrow (dS_i) \leftarrow (d\vec{v}_a)$.

As in Section 6 of paper II, we denote the added edge as AB , and the four added faces will be ABC , ABD , ABE and ABF . We append the derivatives of areas of the first three faces to the minor of matrix $(\partial S_i/\partial \vec{v}_a)$, while the derivatives w.r.t. S_{ABF} — to the minor of matrix $(\partial \Omega_a/\partial S_i)$. We denote those minors simply as $\text{minor}(\partial S_i/\partial \vec{v}_a)$, etc.: we consider only a single minor for every matrix, and there is no risk of confusion.

As concerns the 4-simplices, we continue to use the notations of paper II for them as well. Namely, under the move $2 \rightarrow 4$, two adjacent 4-simplices $ACDEF = \hat{B}$ and $BCDFE = -\hat{A}$ are replaced with four ones: $ABCDE = \hat{F}$, $ABCFD = -\hat{E}$, $ABCEF = \hat{D}$ and $ABDFE = -\hat{C}$.

Formula (II.10) shows that under the move $2 \rightarrow 4$ the important for us minor of matrix $(\partial S_i/\partial \vec{v}_a)$ gets multiplied by

$$\frac{3 V_{\hat{F}} l_{AB}^5}{S_{ABC} S_{ABD} S_{ABE}}. \quad (8)$$

As concerns the minor of matrix $(\partial \Omega_a/\partial S_i)$, it gets multiplied, according to formula (II.15), by

$$\frac{S_{ABF}}{24} \frac{V_{\hat{A}} V_{\hat{B}}}{V_{\hat{C}} V_{\hat{D}} V_{\hat{E}}}. \quad (9)$$

It can be easily derived from here that the following value is conserved under moves $2 \rightarrow 4$:

$$\frac{\text{minor} \left(\frac{\partial \Omega_a}{\partial S_i} \right)}{\text{minor} \left(\frac{\partial S_i}{\partial \vec{v}_a} \right)} \frac{\prod_{\substack{\text{over all} \\ \text{4-simplices}}} V \prod_{\substack{\text{over all} \\ \text{edges}}} 72 l^5}{\prod_{\substack{\text{over all} \\ \text{2-dim. faces}}} S}. \quad (10)$$

Comparing this with formula (7), we see that the value (10) is conserved under moves $3 \rightarrow 3$ as well (of course, $\text{minor}(\partial \Omega_a/\partial S_i)$ is the very same thing as $\det \mathcal{B}$ in formula (7)). Besides, it is again evident that the acyclicity property is conserved.

7 How the algebraic complex changes under moves

$$1 \leftrightarrow 5$$

Like in the previous Section, we consider, for concreteness, only a move in one direction, namely $1 \rightarrow 5$. Under such move, a new *vertex* is added to the simplicial complex — denote it F — which brings about the decomposition of one 4-simplex — denote it $ABCDE$ — into five 4-simplices, which we denote in the style of the previous Section as $\hat{A}$, $\hat{B}$, $\hat{C}$, $\hat{D}$ and $\hat{E}$.

We will explain how we extend the minors that enter in the torsion of complex (5) or (6), starting from the left, that is from the mapping $(dx \text{ and } dg) \rightarrow (dL_a)$ (the minor of the “preceding” mapping $\mathfrak{a} \rightarrow (dx \text{ and } dg)$ does not change, see the remark

after formula (12)). The length of columns of coordinate differentials dx increases by 4: dx_F , dy_F , dz_F and dt_F are added to them — the differentials of four coordinates of the new vertex F . Hence, we must choose four dL , corresponding to four new rows of the minor. Let those be dL_{AF} , dL_{BF} , dL_{CF} and dL_{DF} . The following formula holds which can be proved by a direct calculation using some easy trigonometry (as well as its analogue in 3 dimensions, see [3, formulas (31) and (32)]):

$$dx_F \wedge dy_F \wedge dz_F \wedge dt_F = \frac{dL_{AF} \wedge dL_{BF} \wedge dL_{CF} \wedge dL_{DF}}{384 V_{\hat{E}}}. \quad (11)$$

Strictly speaking, here a $\pm$ sign should have been added, but we take interest in equalities of such kind *to within their sign* (as well as in the preceding Section).

It follows from formula (11) that the important for us *minor of the matrix of mapping* (dx and dg) $\rightarrow$ (dL_a) *gets multiplied by*

$$384 V_{\hat{E}}. \quad (12)$$

Note also that the parameters dg responsible for continuous deformations of representation f are “used up” on algebra $\mathfrak{a}$ (if they existed at all) in the sense that their corresponding rows are included in another minor, that of mapping $\mathfrak{a} \rightarrow (dx \text{ and } dg)$, which does not change under our move.

Now consider the mapping $(dL_a) \rightarrow (d\omega_i)$. Here, only one column is added to the minor, corresponding to the still “available” dL , namely dL_{EF} . On the side of space $(d\omega_i)$, we add the row corresponding to $d\omega_{DEF}$. In consequence, the *minor of the matrix of mapping* $(dL_a) \rightarrow (d\omega_i)$ *gets multiplied by*

$$\frac{S_{DEF}}{24} \frac{V_{\hat{E}} V_{\hat{F}}}{V_{\hat{A}} V_{\hat{B}} V_{\hat{C}}} \quad (13)$$

(cf. formula (9)).

Now we switch from sequence (6) to sequence (5). Nine rows and columns must be added to the minor of mapping $(d\vec{v}_a) \rightarrow (dS_i)$. The columns correspond to the nine “still free” dS_i , that is dS_i for all added faces i except $i = DEF$. We are going to consider them all, step by step, choosing for them on our way nine components of vectors $d\vec{v}_a$ (while their total number is 15: five edges, each having three deviation components).

Matrix $(\partial S_i / \partial \vec{v}_a)$ contains many zeroes (the same applies, by the way, to the other matrices with which we are occupied here): nonzero entries appear only where edge a enters in the boundary of face i . In consequence, many of its submatrices have a block-triangular form so that their determinants (i.e., minors of $(\partial S_i / \partial \vec{v}_a)$) factorize. In particular, it will be convenient for us to include the derivatives w.r.t. *all* components of deviations of edges DF and EF in the minor with which we are occupied now: we will find out that some *separate factors* correspond to them as their contributions to the quantity by which our minor is multiplied. These factors are

$$\frac{dS_{ADF} \wedge dS_{BDF} \wedge dS_{CDF}}{(dv_{DF})_x \wedge (dv_{DF})_y \wedge (dv_{DF})_z} = \frac{3 V_{\hat{E}} l_{DF}^5}{S_{ADF} S_{BDF} S_{CDF}} \quad (14)$$

(cf. formula (8)) for edge DF and

$$\frac{dS_{AEF} \wedge dS_{BEF} \wedge dS_{CEF}}{(dv_{EF})_x \wedge (dv_{EF})_y \wedge (dv_{EF})_z} = \frac{3 V_{\hat{D}} l_{EF}^5}{S_{AEF} S_{BEF} S_{CEF}} \quad (15)$$

for edge EF .

There remain three area differentials — dS_{ABF} , dS_{ACF} and dS_{BCF} — for which we must choose three from nine components of vectors $d\vec{v}_{AF}$, $d\vec{v}_{BF}$ and $d\vec{v}_{CF}$. To this end, we first choose three axes for *each* of these vectors, in order to take their projections onto these axes as their components. Certainly, our triples of axes will make up orthonormal coordinate systems in the three-dimensional spaces orthogonal to corresponding edges. Some easy reasoning shows that the torsion of complex does not depend on a specific choice of such coordinate systems.

We require that the x axes (different!) for $d\vec{v}_{AF}$ and $d\vec{v}_{BF}$ lie in the plane ABF (this fixes their directions, because they must also be orthogonal to the respective edges). The y axis (common) for $d\vec{v}_{AF}$ and $d\vec{v}_{BF}$ will lie in the three-dimensional hyperplane $ABCF$ (and be, of course, orthogonal to plane ABF). The z axis — *common for all three* $d\vec{v}_{AF}$, $d\vec{v}_{BF}$ and $d\vec{v}_{CF}$ — will be orthogonal to hyperplane $ABCF$.

It remains to choose the directions of axes x and y for $d\vec{v}_{CF}$. It is enough to fix the direction of the x axis: we choose it to be orthogonal to plane BCF .

Now we choose three components of vectors $d\vec{v}$, in order to include the derivatives with respect to them into the minor of matrix $(\partial S_i / \partial \vec{v}_a)$. These will be $(dv_{BF})_x$, $(dv_{CF})_x$ and $(dv_{CF})_y$.

The determinant

$$\frac{dS_{ABF} \wedge dS_{ACF} \wedge dS_{BCF}}{(dv_{BF})_x \wedge (dv_{CF})_x \wedge (dv_{CF})_y} \quad (16)$$

again factorizes: it equals the product of the quantity $\partial S_{ABF} / (\partial v_{BF})_x = l_{BF}$ by the determinant $\frac{dS_{ACF} \wedge dS_{BCF}}{(dv_{CF})_x \wedge (dv_{CF})_y}$. But this determinant, too, factorizes due to the fact that dS_{BCF} does not depend on $(dv_{CF})_x$ (recall how we chose the direction of axis x for this vector). Finally we find that expression (16) equals

$$l_{BF} l_{CF}^2 \sin \alpha, \quad (17)$$

where α is the angle between planes ACF and BCF .

Here is the expression — the product of expressions (14), (15) and (17) — by which the minor of matrix $(\partial S_i / \partial \vec{v}_a)$ is multiplied as a result of the move $1 \rightarrow 5$:

$$\frac{9 V_{\hat{E}} V_{\hat{D}} l_{DF}^5 l_{EF}^5}{S_{ADF} S_{BDF} S_{CDF} S_{AEF} S_{BEF} S_{CEF}} l_{BF} l_{CF}^2 \sin \alpha. \quad (18)$$

It remains to consider one more minor that changes under the move $1 \rightarrow 5$ — the minor of matrix $(\partial \vec{v}_a / \partial \sigma)$. It involves the components $(dv_{AF})_x$, $(dv_{AF})_y$, $(dv_{AF})_z$, $(dv_{BF})_y$, $(dv_{BF})_z$ and $(dv_{CF})_z$ of edge deviations and, on the other hand, all six components of bivector $d\sigma$ corresponding to vertex F .

Bivector $d\sigma$ can be thought of as an element of Lie algebra $\mathfrak{so}(4)$, and its components — as infinitesimal rotation angles within the six coordinate planes. We have to choose a proper coordinate system for it. This time, we will denote its axes by numbers 1, 2, 3, 4 (rather than letters x, y, z, t).

We choose axes 1 and 2 to lie in the plane ABF : axis 1 will be orthogonal to vector $\overrightarrow{BF}$, and axis 2 — parallel to it. Axis 3 will lie in the hyperplane $ABCF$ (and, of course, orthogonal to ABF), and axis 4 — orthogonal to $ABCF$.

Now (the thing we are already accustomed to) the value by which the minor is multiplied again factorizes. First (cf. formula (2)), the components $(dv_{AF})_z$, $(dv_{BF})_z$ and $(dv_{CF})_z$ are determined just by three rotation angles in the direction of axis 4, i.e. $d\sigma_{14}$, $d\sigma_{24}$ and $d\sigma_{34}$. Namely, they are connected by means of the following 3×3 submatrix of matrix $(\partial \vec{v}_a / \partial \sigma)$ (see again (2)):

$$\begin{pmatrix} \frac{(AF)_1}{L_{AF}} & \frac{(AF)_2}{L_{AF}} & \frac{(AF)_3}{L_{AF}} \\ \frac{(BF)_1}{L_{BF}} & \frac{(BF)_2}{L_{BF}} & \frac{(BF)_3}{L_{BF}} \\ \frac{(CF)_1}{L_{CF}} & \frac{(CF)_2}{L_{CF}} & \frac{(CF)_3}{L_{CF}} \end{pmatrix}, \quad (19)$$

where, for instance, $(AF)_1$ is the component of vector $\overrightarrow{AF}$ along axis 1. The determinant of matrix (19) is

$$\frac{6 V_{ABCF}}{L_{AF} L_{BF} L_{CF}} = \frac{4 S_{ACF} S_{BCF} \sin \alpha}{L_{AF} L_{BF} l_{CF}^3}, \quad (20)$$

where V_{ABCF} is the *three-dimensional* volume, while α is the same angle as in formula (17).

There remain three rotations within hyperplane $ABCF$, and three components $(dv_{AF})_x$, $(dv_{AF})_y$ and $(dv_{BF})_y$. Here, too, the factorizability persists: the rotation within the plane ABF affects only $(dv_{AF})_x$, and this gives the factor

$$\frac{(\partial v_{AF})_x}{\partial \sigma_{12}} = \frac{1}{l_{AF}}. \quad (21)$$

The rotation within the plane orthogonal to BF affects only $(dv_{AF})_y$, which gives the factor

$$\frac{(\partial v_{AF})_y}{\partial \sigma_{13}} = \frac{\sin \beta}{l_{AF}}, \quad (22)$$

β being the angle between edges AF and BF . Now the last rotation remains with the corresponding factor

$$\frac{(\partial v_{BF})_y}{\partial \sigma_{23}} = \frac{1}{l_{BF}}. \quad (23)$$

Taking the product of the right-hand sides of formulas (20), (21), (22) and (23), we find the factor by which the minor of matrix $(\partial \vec{v}_a / \partial \sigma)$ gets multiplied under our move $1 \rightarrow 5$:

$$\frac{8 S_{ACF} S_{BCF} S_{ABF}}{l_{AF}^5 l_{BF}^4 l_{CF}^3} \cdot \sin \alpha, \quad (24)$$

where we have taken into account that $\sin \beta = 2 S_{ABF} / l_{AF} l_{BF}$.

In the same way as in Subsections 6.1 and 6.2, it is clear from the performed reasoning and calculations that the property of acyclicity of the complex is conserved under the considered moves.

8 The formula for the manifold invariant

In the two preceding Sections we have analyzed the behaviour of the minors whose alternated product makes up the torsion of the complex (5) or (6) (if this complex is acyclic), under Pachner moves $3 \rightarrow 3$, $2 \leftrightarrow 4$ and $1 \leftrightarrow 5$. We have now to unite these results to obtain the quantity which *does not depend on a triangulation*, i.e. an *invariant of a four-dimensional piecewise-linear manifold*.

We choose the signs in the alternated product in such way that our formula for the invariant look as similar as possible to the “three-dimensional” formula (II.5). Namely, we take the minor of the matrix of mapping $\mathfrak{a} \rightarrow (dx \text{ and } dg)$ raised in the power -1 , then the minor of the matrix of mapping $(dx \text{ and } dg) \rightarrow (dL_a)$ raised in the power $+1$ and so on. We will get the factor by which the so defined torsion τ is multiplied under the move $1 \rightarrow 5$ when we multiply the expressions (12) and (18) and divide by the product of (13) and (24). The result can be written as

$$2^7 \cdot 3^4 \cdot \frac{V_{\hat{A}} V_{\hat{B}} V_{\hat{C}} V_{\hat{D}} V_{\hat{E}}}{V_{\hat{F}}} \cdot \frac{\prod_{\substack{\text{over new} \\ \text{edges}}} l^5}{\prod_{\substack{\text{over new} \\ \text{2-dim. faces}}} S}.$$

Comparing this with the results of Section 6 (formulas (7) and (10)), we find the following final expression for the invariant of a four-dimensional manifold in terms of the torsion of complex (5) or (6) and other Euclidean geometric values:

$$I = 2^{-16} \cdot 3^{-12} \cdot \frac{\tau \cdot \prod_{\substack{\text{over all} \\ \text{2-dim. faces}}} S}{\prod_{\substack{\text{over all} \\ \text{4-simplices}}} V \prod_{\substack{\text{over all} \\ \text{edges}}} 72 l^5} \cdot (2^8 \cdot 3^6)^{(\text{number of vertices})}. \quad (25)$$

The factor $2^{-16} \cdot 3^{-12}$ has been added in order that the invariant be equal to unity for the sphere S^4 , see the following Section.

9 Example: sphere S^4

The fundamental group of sphere S^4 is trivial, thus algebra $\mathfrak{a}$ in sequence (6) will be the whole Lie algebra $\mathfrak{e}_4$ of motions of the four-dimensional Euclidean space. There are

obviously no continuous parameters dg describing the deformations of representation $\pi_1(S^4) \rightarrow E_4$. The space of “trivial” vertex deviations is ten-dimensional and consists of deviations of type (4) with arbitrary $d\tau_{\alpha\beta}$ and ds_β .

We take the canonical triangulation of S^4 consisting of two 4-simplices, both with vertices A, B, C, D and E . For such triangulation, all deficit angles $d\omega_i$ and $d\Omega_a$ are identically equal to zero, because any ω_i is obtained by summing two terms differing only in sign (the $+$ sign is ascribed to one of 4-simplices, while the $-$ sign to the other one, see I, Section 3). Thus, matrix $(\partial\Omega_a/\partial S_i)$ and its conjugate $(\partial\omega_i/\partial L_a)$ are zero. This means that the torsion of complex (5) or (6) factorizes in the product over two following sequences:

$$0 \rightarrow \mathfrak{e}_4 \rightarrow (dx) \rightarrow (dL_a) \rightarrow 0 \quad (26)$$

and

$$0 \leftarrow (dS_i) \leftarrow (d\vec{v}_a) \leftarrow (d\sigma) \leftarrow (d\sigma)_0 \leftarrow 0. \quad (27)$$

We fix a Euclidean coordinate system in the space $\mathbb{R}_4$, with axes x, y, z, t . We place the vertex A of our triangulation into the origin of coordinates, and the remaining vertices — in points $B(1, 0, 0, 0)$, $C(0, 1, 0, 0)$, $D(0, 0, 1, 0)$ and $E(0, 0, 0, 1)$.

9.1 Calculations for sequence (26)

The ten-dimensional algebra $\mathfrak{e}_4$ consists of infinitesimal rotations and translations (of course, we measure the rotations within the six coordinate planes in radians, and the translations — in the units of coordinate axes), whereas the twenty-dimensional

space (dx) — of column vectors $\begin{pmatrix} dx_A \\ \vdots \\ dt_E \end{pmatrix}$. We take the minor of matrix of mapping $\mathfrak{e}_4 \rightarrow (dx)$ corresponding to the following *ten* coordinate differentials: $dx_A, dy_A, dz_A, dt_A, dy_B, dz_B, dt_B, dz_C, dt_C, dt_D$.

From this minor, a unity factor splits off at once corresponding to the mapping (translations) $\rightarrow$ (vertex A coordinates); this splitting off is caused by the fact that *rotations do not affect* the coordinates of A . There remains the mapping of six rotations into sets $(dy_B, dz_B, dt_B, dz_C, dt_C, dt_D)$. Factorization still works here: rotations within the planes xy , xz and xt affect only point B (from those remaining) and so on. The result is: *the minor of mapping $\mathfrak{e}_4 \rightarrow (dx)$ equals unity* (as usual, to within a sign).

Now we consider the minor of matrix of the mapping $(dx) \rightarrow (dL_a)$ corresponding to the set $(dx_B, dx_C, dy_C, dx_D, dy_D, dz_D, dx_E, dy_E, dz_E, dt_E)$. It, too, factorizes into the product of determinants of mappings

$$(dx_E, dy_E, dz_E, dt_E) \rightarrow (dL_{AE}, dL_{BE}, dL_{CE}, dL_{DE}), \quad (28)$$

$$(dx_D, dy_D, dz_D) \rightarrow (dL_{AD}, dL_{BD}, dL_{CD}), \quad (29)$$

$$(dx_C, dy_C) \rightarrow (dL_{AC}, dL_{BC}), \quad (30)$$

$$dx_B \rightarrow dL_{AB}. \quad (31)$$

Recall that L is the squared length of a corresponding edge. A direct calculation shows that the determinants of mappings (28), (29), (30) and (31) equal 2^4 , 2^3 , 2^2 and 2 , respectively.

Conclusion: the multiplicative contribution of sequence (26) to the torsion is 2^{10} .

9.2 Calculations for sequence (27)

We start from the right, i.e. from the mapping $(d\sigma)_0 \rightarrow (d\sigma)$. The ten-dimensional space $(d\sigma)_0$ consists of the components of antisymmetric tensor $d\tau_{\alpha\beta}$ and vector ds_β , whereas the thirty-dimensional space $(d\sigma)$ — of the components of all five vertices' deviations. The mapping is given by formula (4).

For composing the minor of matrix of the mapping $(d\sigma)_0 \rightarrow (d\sigma)$, we choose the following ten components of tensors $d\sigma_A, \dots, d\sigma_E$: we take *all six* components of $d\sigma_A$ and *one* component for each of the remaining deviations, namely, $(d\sigma_B)_{zt}$, $(d\sigma_C)_{zt}$, $(d\sigma_D)_{xy}$ and $(d\sigma_E)_{xy}$. It makes sense to remind here that this choice depends on us (the only requirement is that minors be nonzero), and this exactly choice was motivated by the convenience of further calculations.

The vector ds_β does not affect $d\sigma_A$, because the radius vector of point A is zero. Hence, our minor factorizes, and the factor corresponding to mapping $d\tau \rightarrow d\sigma_A$ is unity (because this is an identical mapping: $d\sigma_A = d\tau$). The remaining 4×4 minor factorizes into the product of four unit factors, because, for instance, $(d\sigma_B)_{zt}$ depends on ds_y only (this can be seen from formula (4) if we replace in it the subscript A with B , ignore $d\tau_{\alpha\beta}$ and observe that the single nonzero component of vector $\vec{r}_B$ is $(r_B)_x$) and so on.

Thus, our selected minor of matrix of the mapping $(d\sigma)_0 \rightarrow (d\sigma)$ turned out to equal unity.

We pass on to the mapping $(d\sigma) \rightarrow (d\vec{v}_a)$. Here one should fix at first a three-dimensional basis for each of ten vectors $d\vec{v}_{AB}, \dots, d\vec{v}_{DE}$. It is done most easily for those edges that begin at point A : edge AB lies on the x axis, so we choose as the coordinate axes for $d\vec{v}_{AB}$ the three remaining axes y , z and t ; then we choose axes in a similar way for the deviations of edges AC , AD and AE .

Each of the remaining six edges lies within some coordinate *plane*, and we will treat them in the following way. Edge BC lies in the plane xy ; we choose the following three axes for $d\vec{v}_{BC}$: the bisector of the angle formed by axes x and y , and also axes z and t ; we follow this model in choosing the axes for deviations of edges BD , BE , CD , CE and DE . We will denote the components of vectors of edge deviations along the bisectors of coordinate angles as $(dv_{BC})_{\text{bis}}$, etc. (keeping in mind that every deviation has its own bisector for an axis).

We must choose twenty components of vectors of edge deviations (because we have twenty components of vertex deviations not included in our previous minor). We choose *all* components for edges AB , AC , AD and AE , and also the following eight components: $(dv_{BD})_y$, $(dv_{BD})_t$, $(dv_{BE})_y$, $(dv_{BE})_z$, $(dv_{CD})_x$, $(dv_{CD})_t$, $(dv_{CE})_x$ and $(dv_{CE})_z$. We drew in Figure 2 the four edges to which these components belong, and wrote out




Figure 2:

these components near the edges (for example, we wrote dv_y and dv_t near the edge BD meaning $(dv_{BD})_y$ and $(dv_{BD})_t$).

Of course, our 20×20 minor again greatly factorizes. For example, the minor corresponding, on the one hand, to the components $(d\sigma_B)_{xy}$, $(d\sigma_B)_{xz}$ and $(d\sigma_B)_{xt}$, and on the other hand — to the three components of $d\vec{v}_{AB}$, factors out (this is because all the components of $d\sigma_A$ are already used up, whereas only they of all the remaining components of $d\sigma$ could influence $d\vec{v}_{AB}$). From formulas of type (2) we can see that this factor equals unity. Similarly, three more minors factor out which are obtained from the above minor by changing $B \rightarrow C$, $x \leftrightarrow y$, or by changing $B \rightarrow D$, $x \leftrightarrow z$, or by changing $B \rightarrow E$, $x \leftrightarrow t$.

After this, there remain the eight already mentioned components of vectors $d\vec{v}$ depicted in Figure 2, and the eight components of $d\sigma$ also depicted in Figure 2, with the understanding that if, e.g., $d\sigma_{yz}$ and $d\sigma_{yt}$ are drawn near the vertex B , then they are $(d\sigma_B)_{yz}$ and $(d\sigma_B)_{yt}$. The corresponding minor factorizes in eight separate factors. The reason for this is that each component of $d\vec{v}$ depends on *only one* component of $d\sigma$. These dependencies are shown by arrows in Figure 2. Note also that the small letters near the vertices denote the axes where these vertices lie.

We see from formulas of type (2) or (3) that all the eight factors equal $\frac{1}{2}$. Consequently, their contribution to the torsion is 2^8 .

It remains for us to consider the mapping $(d\vec{v}_a) \rightarrow (dS_i)$. As we remember, all the components of vectors $d\vec{v}$ corresponding to edges AB , AC , AD and AE are already used up. For the remaining six edges, we have the components of $d\vec{v}$ along the bysectors of coordinate angles, as well as $(dv_{BC})_z$, $(dv_{BC})_t$, $(dv_{DE})_x$ and $(dv_{DE})_y$ (these latter belong to the two edges *absent* from Figure 2).

Each of the six differentials dS_i , where face i contains vertex A , is influenced by only one component of only one of the remaining vectors $d\vec{v}$. In this way, six factors

appear: $\partial S_{ABC}/(\partial v_{BC})_{\text{bis}} = \sqrt{2}$ and five other ones, all equal to it. So, here we have the contribution to the torsion equal to 2^3 .

There remains the minor whose rows correspond to components dS_{BCD} , dS_{BCE} , dS_{BDE} and dS_{CDE} , while columns — to $(dv_{BC})_z$, $(dv_{BC})_t$, $(dv_{DE})_x$ and $(dv_{DE})_y$. Here, too, each component dS turns out to depend on only one component dv . We get the product of four partial derivatives, all equal to each other; for example, one of them is

$$\frac{\partial S_{BCD}}{(\partial v_{BC})_z} = l_{BC} \cdot \cos \gamma = \sqrt{2} \cdot \sqrt{\frac{2}{3}} = \frac{2}{\sqrt{3}},$$

where γ is the angle between face BCD and axis z . Thus, here the contribution to the torsion is $2^4 \cdot 3^{-2}$.

Conclusion: sequence (27) makes the multiplicative contribution to the torsion, equal to $2^8 \cdot 2^3 \cdot 2^4 \cdot 3^{-2} = 2^{15} \cdot 3^{-2}$.

9.3 The result: invariant for sphere S^4

Combining the conclusions made in the end of two previous subsections, we find that the torsion for sphere S^4 is

$$\tau(S^4) = 2^{25} \cdot 3^{-2}.$$

Now we calculate the products entering in formula (25).

In our complex, there are six two-dimensional faces of area $1/2$ and four faces of area $\sqrt{3}/2$. Next, $\prod S = 3^2/2^{10}$. Then, there are two 4-simplices, both of volume $1/24$, thus $\prod V = 2^{-6} \cdot 3^{-2}$. Finally, there are four edges of length 1 and six edges of length $\sqrt{2}$, thus $\prod 72 l^5 = 2^{45} \cdot 3^{20}$.

All this together leads to the formula announced in the end of Section 8:

$$I(S^4) = 1. \tag{32}$$

10 Discussion

So, in the case of sphere S^4 the complex turned out to be acyclic, and we managed to calculate its torsion (and our invariant). The largest determinant that we had to deal with was of sizes 20×20 , but, luckily, it factorized in a product of smaller determinants.

Hopefully, new properties of our invariants will be discovered with time, which will simplify the calculations, and some relevant techniques will be elaborated. This will give the real possibility to calculate the invariants for a large enough manifold zoo. At this moment, one of the interesting questions is what we will get for the product of two-dimensional spheres $S^2 \times S^2$ and whether we will be able to do something if the corresponding complex turns out *not* to be acyclic.

One more problem is the generalization of our complexes and finding their possible quantum analogues. One can begin with constructing a complex based on the $SL(2)$ -solution to the pentagon equation from paper [5].

It looks quite plausible that our constructions can be generalized in such way that they include also the *Reidemeister torsion*. In prospect, one can think about the creation of a new general theory that will combine the ideas of the algebra of acyclic complexes and quantum topology.

Acknowledgements. The work has been performed with the partial financial support from Russian Foundation for Basic Research, Grant no. 01-01-00059.

References

- [1] *I. G. Korepanov*. Euclidean 4-simplices and invariants of four-dimensional manifolds: I. Moves $3 \rightarrow 3$. Theor. Math. Phys., Volume 131 (2002), 765–774.
- [2] *I. G. Korepanov*. Euclidean 4-simplices and invariants of four-dimensional manifolds: II. An algebraic complex and moves $2 \leftrightarrow 4$. Theor. Math. Phys., Volume 133 (2002), 1338–1347.
- [3] *I. G. Korepanov*. Invariants of PL manifolds from metrized simplicial complexes. Three-dimensional case. J. Nonlin. Math. Phys., Volume 8, number 2 (2001), 196–210.
- [4] *I. G. Korepanov, E. V. Martyshev*. Distinguishing three-dimensional lens spaces $L(7, 1)$ and $L(7, 2)$ by means of classical pentagon equation. J. Nonlin. Math. Phys., Volume 9, number 1 (2002), 86–98.
- [5] *I. G. Korepanov, E. V. Martyshev*. A Classical Solution of the Pentagon Equation Related to the Group $SL(2)$. Theor. Math. Phys., Volume 129 (2001), 1320–1324.